

— Summary of Curriculum Vitae — (April, 2016)



Name: Kiyonori KURIKI
Position: Associate Professor
Degrees awarded:
PhD: Nagoya City University, 2002
MSc: Meijyo University, 1996

Work address (2009 - -> Current):

Laboratory of Public Health, Department of Nutritional Sciences, School of Food and Nutritional Sciences, and Graduate Program in Food and Nutritional Sciences, Graduate School of Integrated Pharmaceutical and Nutritional Sciences, University of Shizuoka

52-1 Yada, Suruga-ku, Shizuoka, 422-8526, Japan

Phone (& FAX): +81 -54 -264 -5563

e-mail: kuriki@u-shizuoka-ken.ac.jp

<http://sfns.u-shizuoka-ken.ac.jp/sfnseng/teachers/page017.html> (English version)

<http://eng.u-shizuoka-ken.ac.jp/> (English version)

Education:

1991-1994: Meijyo University, Faculty of Pharmacy

1995-1996: Meijyo University Graduate School

1998-2002: Department of Health and Preventive Medicine,
Nagoya City University Graduate School of Medical Sciences,

Program Experience:

2002-2005: Research Fellowships of the Japan Society for the Promotion of Science

Division of Epidemiology and Prevention, Aichi Cancer Center Research Institute

2005-2007: Awardees of Research Resident Fellowship from the Foundation for Promotion of Cancer

Research (Japan) for the 3rd Term Comprehensive 10- Year Strategy for Cancer Control

(Division of Epidemiology and Prevention, Aichi Cancer Center Research Institute)

2008: INSERM, E3N-EPIC, ERI20 Equipe Nutrition, Hormones et Cancer,

Institut de cancérologie Gustave Roussy, FRANCE

Chercheur Etranger sur Poste Vert (Foreigner Research Fellowships: PD)

* Institut national de la santé et la recherche médicale (INSERM)

List of published papers (*: Corresponding author)

Original articles

49. Endoh K, Kuriki K (*), Kasezawa N, Tohyama K, Goda T. Interactions between self-reported psychological stress and habitual exercise on the dietary intake of Japanese men and women: a large-scale cross-sectional study. *Asian Pac J Cancer Prev.* (in press)
48. Shirai Y, Kuriki K (*), Endoh K, Miyauchi R, Kasezawa N, Tohyama K, Goda T. Positive linear dose-response relationships, but no J-shaped relationship, between drinking habits and estimated glomerular filtration rate in middle-aged Japanese men. *Alcohol.* 2016, 51, 71-77.
47. Yamada N, Shibata K, Fuku M, Kuriki K, Goto C, Tokudome Y, Miyata M, Maeda K, Hoshino H, Suzuki S, Marumoto M, Ichikawa H, Agawa H, Ichikawa N, Arakawa K, Hosono A, Kobayashi M, Tokudome S. Changes of tryptophan metabolism in Japanese runners during an ultra-marathon race. *Sport Sci Health*, DOI 10.1007/s11332-016-0257-1.
46. Hisada T, Endoh K, Kuriki K (*). Inter- and intra-individual variations in seasonal and daily stabilities of the human gut microbiota in Japanese. *Arch Microbiol*, 2015, 197, 919-34.
45. Endoh K, Kuriki K (*), Kasezawa N, Tohyama K, Goda T. Association between smoking status and food and nutrient consumption in Japanese: a large-scale cross-sectional study. *Asian Pac J Cancer Prev*, 2015, 16, 6527-34.
44. Endoh K, Kuriki K (*), Kasezawa N, Tohyama K, Goda T. Interactions between psychological stress and drinking status in relation to diet among middle-aged men and women: large-scale cross-sectional study in Japan. *J. Nutr. Sci. Vitaminol (Tokyo)*, 2015, 61, 64-72.
43. Tokudome S, Kuriki K, Yokoyama Y, Sasaki M, Joh T, Kamiya T, Cheng J, Ogawa K, Shirai T, Imaeda N, Goto C, Tokudome Y, Ichikawa H, Okuyama H. Dietary n-3/long-chain n-3 polyunsaturated fatty acids for prevention of sporadic colorectal tumors: A randomized controlled trial in polypectomized participants. *Prostaglandins, Leukotrienes and Essential Fatty Acids*, 2015;94:1-11.
42. Imaeda N, Kuriki K, Fujiwara N, Goto C, Tokudome Y, Tokudome S. Usual dietary intakes of selected trace elements (Zn, Cu, Mn, I, Se, Cr, and Mo) and biotin revealed by a survey of four-season 7-consecutive day weighed dietary records in middle-aged Japanese dietitians. *Nutr Sci Vitaminol (Tokyo)*. 2013;59:281-8.
41. Mochizuki K, Misaki Y, Miyauchi R, Takabe S, Shimada M, Kuriki K, Ichikawa Y, Goda T. A higher rate of eating is associated with higher circulating interleukin-1 β concentrations in Japanese men not being treated for metabolic diseases. *Nutrition*, 2012;28:978-83.
40. Wang J, Jiang J, Zhao Y, Gajalakshmi V, Kuriki K, Suzuki S, Nagaya T, Nakamura S, Akasaka S, Ishikawa H, Tokudome S. Genetic polymorphisms of glutathione S-transferase genes and susceptibility to colorectal cancer: A case-control study in an Indian population. *Cancer Epidemiol.* 2011;35:66-72
39. Wang J, Zhao Y, Jiang J, Gajalakshmi V, Kuriki K, Nakamura S, Akasaka S, Ishikawa H, Suzuki S, Nagaya T, Tokudome S. Polymorphisms in DNA repair genes XRCC1, XRCC3 and XPD, and colorectal cancer risk: a case-control study in an Indian population. *J Cancer Res Clin Oncol.*

2010;136:1517-25.

38. Maillard V, Kuriki K, Lefebvre B, Boutron-Ruault MC, Lenoir GM, Joulin V, Clavel-Chapelon F, Chajès V. Serum carotenoid, tocopherol and retinol concentrations and breast cancer risk in the E3N-EPIC Study. *Int J Cancer*. 2010;127:1188-96.
37. Hattori N, Hayashi T, Nakachi K, Ichikawa H, Goto C, Tokudome Y, Kuriki K, Hoshino H, Shibata K, Yamada N, Fuku M, Suzuki S, Nagaya T, Kobayashi M, Tokudome S. Changes in Concentrations of Serum Reactive Oxygen Species during a Two-day Ultra-marathon Race in Japanese Male Amateur Runners. *Int J Sports Med* 2009;30:426-9.
36. Xiang J, Nagaya T, Huang XE, Kuriki K, Imaeda N, Tokudome Y, Sato J, Fujiwara N, Maki S, Tokudome S. Sex and Seasonal Variations of Plasma Retinol, α -Tocopherol, and Carotenoid Concentrations in Japanese Dietitians. *Asian Pac J Cancer Prev*. 2008;9:413-6.
35. Kuriki K (*), Mutoh M, Tajima K, Wakabayashi Keiji, Tatematsu M. Relationships between intestinal polyp formation and fatty acid levels in plasma, erythrocytes and intestinal polyps in Min mice. *Cancer Sci*. 2008;99:2410-6.
34. Ghadimi R, Kuriki K, Tsuge S, Takeda E, Imaeda N, Suzuki S, Sawai A, Takekuma K, Hosono A, Tokudome Y, Goto C, Esfandiary I, Nomura H, Tokudome S. Serum concentrations of Fatty acids and colorectal adenoma risk: a case-control study in Japan. *Asian Pac J Cancer Prev*. 2008;9:111-8.
33. Yoshida S, Okazaki Y, Yamashita T, Ueda H, Ghadimi R, Hosono A, Tanaka T, Kuriki K, Suzuki S, Tokudome S. Analysis of human oral mucosa ex vivo for fatty acid compositions using Fourier-transform infrared spectroscopy. *Lipids*. 2008;43:361-72.
32. Miyata M, Kasai H, Kawai K, Yamada N, Tokudome M, Ichikawa H, Goto C, Tokudome Y, Kuriki K, Hoshino H, Shibata K, Suzuki S, Kobayashi M, Goto H, Ikeda M, Otsuka T, Tokudome S. Changes of urinary 8-hydroxydeoxyguanosine levels during a two-day ultramarathon race period in Japanese non-professional runners. *Int J Sports Med*. 2008;29:27-33.
31. Kuriki K (*), Wakai K, Matsuo K, Hiraki A, Suzuki T, Yamamura Y, Yamao K, Nakamura T, Tatematsu M, Tajima K. Gastric cancer risk and erythrocyte composition of docosahexaenoic acid with anti-inflammatory effects. *Cancer Epidemiol Biomarkers Prev*. 2007;16:2406-15.
30. Kuriki K (*), Hirose K, Wakai K, Matsuo K, Ito H, Suzuki T, Hiraki A, Saito T, Iwata H, Tatematsu M, Tajima K. Breast cancer risk and erythrocyte compositions of n-3 highly unsaturated fatty acids in Japanese. *Int J Cancer*. 2007;121:377-85.
29. Kuriki K (*), Hirose K, Tajima K. Diabetes and cancer risk for all and specific sites among Japanese men and women. *Eur J Cancer Prev*. 2007;16:83-9.
28. Kuriki K (*), Wakai K, Hirose K, Matsuo K, Ito H, Suzuki T, Saito T, Kanemitsu Y, Hirai T, Kato T, Tatematsu M, Tajima K. Risk of colorectal cancer is linked to erythrocyte compositions of fatty acids as biomarkers for dietary intakes of fish, fat, and fatty acids. *Cancer Epidemiol Biomarkers Prev*. 2006;15:1791-8.
27. Kuriki K (*), Tajima K, Tokudome S. Accelerated solvent extraction for quantitative measurement of fatty acids in plasma and erythrocytes. *Lipids*. 2006;41:605-14.

26. Kuriki K (*), Hirose K, Matsuo K, Wakai K, Ito H, Kanemitsu Y, Hirai T, Kato T, Hamajima N, Takezaki T, Suzuki T, Saito T, Tanaka R, Tajima K. Meat, milk, saturated fatty acids, the Pro12Ala and C161T polymorphisms of the PPARgamma gene and colorectal cancer risk in Japanese. *Cancer Sci.* 2006;97:1226-35.
25. Suzuki T, Wakai K, Matsuo K, Hirose K, Ito H, Kuriki K, Sato S, Ueda R, Hasegawa Y, Tajima K. Effect of dietary antioxidants and risk of oral, pharyngeal and laryngeal squamous cell carcinoma according to smoking and drinking habits. *Cancer Sci.* 2006;97:760-7.
24. Wakai K, Hirose K, Matsuo K, Ito H, Kuriki K, Suzuki T, Kato T, Hirai T, Kanemitsu Y, Tajima K. Dietary risk factors for colon and rectal cancers: a comparative case-control study. *J Epidemiol.* 2006;16:125-35.
23. Jiang J, Wang J, Suzuki S, Gajalakshmi V, Kuriki K, Zhao Y, Nakamura S, Akasaka S, Ishikawa H, Tokudome S. Elevated risk of colorectal cancer associated with the AA genotype of the cyclin D1 A870G polymorphism in an Indian population. *J Cancer Res Clin Oncol.* 2006;132:193-9.
22. Wang J, Gajalakshmi V, Jiang J, Kuriki K, Suzuki S, Nagaya T, Nakamura S, Akasaka S, Ishikawa H, Tokudome S. Associations between 5,10-methylenetetrahydrofolate reductase codon 677 and 1298 genetic polymorphisms and environmental factors with reference to susceptibility to colorectal cancer: a case-control study in an Indian population. *Int J Cancer.* 2006;118:991-7.
21. Goto C, Tokudome Y, Imaeda N, Takekuma K, Kuriki K, Igarashi F, Ikeda M, Tokudome S. Validation study of fatty acid consumption assessed with a short food frequency questionnaire against plasma concentration in middle-aged Japanese people. *Scand J Nutr* 2006;50:77-82.
20. Jiang J, Gajalakshmi V, Wang J, Kuriki K, Suzuki S, Nakamura S, Akasaka S, Ishikawa H, Tokudome S. Influence of the C161T but not Pro12Ala polymorphism in the peroxisome proliferator-activated receptor-gamma on colorectal cancer in an Indian population. *Cancer Sci.* 2005;96:507-12.
19. Jiang J, Suzuki S, Xiang J, Kuriki K, Hosono A, Arakawa K, Wang J, Nagaya T, Kojima M, Katsuda N, Tokudome S. Plasma carotenoid, alpha-tocopherol and retinol concentrations and risk of colorectal adenomas: A case-control study in Japan. *Cancer Lett.* 2005;226:133-41.
18. Kuriki K (*), Hamajima N, Chiba H, Kanemitsu Y, Hirai T, Kato T, Saito T, Matsuo K, Koike K, Tokudome S, Tajima K. Increased risk of colorectal cancer due to interactions between meat consumption and the CD36 gene A52C polymorphism among Japanese. *Nutr Cancer.* 2005;51:170-7.
17. Kuriki K (*), Hamajima N, Chiba H, Kanemitsu Y, Hirai T, Kato T, Saito T, Matsuo K, Koike K, Tokudome S, Tajima K. Relation of the CD36 gene A52C polymorphism to the risk of colorectal cancer among Japanese, with reference to with the aldehyde dehydrogenase 2 gene Glu487Lys polymorphism and drinking habit. *Asian Pac J Cancer Prev.* 2005;6 62-8.
16. Tokudome S, Soeripto, Triningsih FX, Ananta I, Suzuki S, Kuriki K, Akasaka S, Kosaka H, Ishikawa H, Azuma T, Moore MA. Rare *Helicobacter pylori* infection as a factor for the very low stomach cancer incidence in Yogyakarta, Indonesia. *Cancer Lett.* 2005;219:57-61.
15. Chiba H, Takezaki T, Neupani D, Kim J, Yoshida S, Mizoguchi E, Takeuchi J, Suzuki J, Tanaka Y, Ito K, Kitamura T, Kuriki K, Wakai K, Samejima K, Sonoda S, Tajima K. An epidemiological study of HBV,

- HCV and HTLV-I in Sherpas of Nepal. *Asian Pac J Cancer Prev.* 2004;5:370-3.
14. Tokudome S, Kuriki K, Yamada N, Ichikawa H, Miyata M, Shibata K, Hoshino H, Tsuge S, Tokudome M, Goto C, Tokudome Y, Kobayashi M, Goto H, Suzuki S, Okamoto Y, Ikeda M, Sato Y. Anthropometric, lifestyle and biomarker assessment of Japanese non-professional ultra-marathon runners. *J Epidemiol.* 2004;14:161-7.
 13. Kuriki K (*), Tokudome S, Tajima K. Association between type II diabetes and colon cancer among Japanese with reference to changes in food intake. *Asian Pac J Cancer Prev.* 2004;5:28-35.
 12. Kuriki K, Nagaya T, Tokudome Y, Imaeda N, Fujiwara N, Sato J, Goto C, Ikeda M, Maki S, Tajima K, Tokudome S. Plasma concentrations of (n-3) highly unsaturated fatty acids are good biomarkers of relative dietary fatty acid intakes: a cross-sectional study. *J Nutr.* 2003;133:3643-50.
 11. Tokudome Y, Kuriki K, Imaeda N, Ikeda M, Nagaya T, Fujiwara N, Sato J, Goto C, Kikuchi S, Maki S, Tokudome S. Seasonal variation in consumption and plasma concentrations of fatty acids in Japanese female dietitians. *Eur J Epidemiol.* 2003;18:945-53.
 10. Wang J, Deng Y, Li L, Kuriki K, Ding J, Pan X, Zhuge X, Jiang J, Luo C, Lin P, Tokudome S. Association of GSTM1, CYP1A1 and CYP2E1 genetic polymorphisms with susceptibility to lung adenocarcinoma: a case-control study in Chinese population. *Cancer Sci.* 2003;94:448-52.
 9. Tokudome S, Nagaya T, Okuyama H, Tokudome Y, Imaeda N, Kitagawa I, Fujiwara N, Ikeda M, Goto C, Ichikawa H, Kuriki K, Takekuma K, Shimoda A, Hirose K, Usui T. Japanese versus Mediterranean Diets and Cancer. *Asian Pac J Cancer Prev.* 2000;1:61-66.
 8. Cheng J, Ogawa K, Kuriki K, Yokoyama Y, Kamiya T, Seno K, Okuyama H, Wang J, Luo C, Fujii T, Ichikawa H, Shirai T, Tokudome S. Increased intake of n-3 polyunsaturated fatty acids elevates the level of apoptosis in the normal sigmoid colon of patients polypectomized for adenomas/tumors. *Cancer Lett.* 2003;10;193:17-24. Erratum in: *Cancer Lett.* 2004;205(2):227.
 7. Tokudome S, Yokoyama Y, Kamiya T, Seno K, Okuyama H, Kuriki K, Cheng J, Nakamura T, Fujii T, Ichikawa H, Itoh M. Rationale and study design of dietary intervention in patients polypectomized for tumors of the colorectum. *Jpn J Clin Oncol.* 2002;32:550-3.
 6. Tokudome Y, Imaeda N, Nagaya T, Ikeda M, Fujiwara N, Sato J, Kuriki K, Kikuchi S, Maki S, Tokudome S. Daily, weekly, seasonal, within- and between-individual variation in nutrient intake according to four season consecutive 7 day weighed diet records in Japanese female dietitians. *J Epidemiol.* 2002;12:85-92.
 5. Kuriki K, Nagaya T, Imaeda N, Tokudome Y, Fujiwara N, Sato J, Ikeda M, Maki S, Tokudome S. Discrepancies in dietary intakes and plasma concentrations of fatty acids according to age among Japanese female dietitians. *Eur J Clin Nutr.* 2002;56:524-31.
 4. Imaeda N, Fujiwara N, Tokudome Y, Ikeda M, Kuriki K, Nagaya T, Sato J, Goto C, Maki S, Tokudome S. Reproducibility of a semi-quantitative food frequency questionnaire in Japanese female dietitians. *J Epidemiol.* 2002;12:45-53.
 3. Hayashi K, Sato J, Fujiwara N, Kajita M, Fukuharu M, Xiaochen HU, Kuriki K, Hoshino H, Kato R, Tokudome S, Sato R. Associations between physical strength, cerebral function and mental health in

independent - living elderly Japanese women. *Environ Health Prev Med* 2002;7:123-8.

2. Tokudome S, Imaeda N, Tokudome Y, Fujiwara N, Nagaya T, Sato J, Kuriki K, Ikeda M, Maki S. Relative validity of a semi-quantitative food frequency questionnaire versus 28 day weighed diet records in Japanese female dietitians. *Eur J Clin Nutr*. 2001;55:735-42.
1. Fukuharu M, Sato J, Ohsawa I, Oshida Y, Kuriki K, Shibata K, Tamagawa T, Nagaya T, Fujiwara N, Tokudome S, Sato Y. Effects of lifestyle factors on ultrasonographically determined bone health in Japanese women. *Public Health*. 2001;115:146-51.

Review

2. Kuriki K (*), Tajima K. The increasing incidence of colorectal cancer and the preventive strategy in Japan. *Asian Pac J Cancer Prev*. 2006;7:495-501.
1. Moore MA, Sobue T, Kuriki K, Tajima K, Tokudome S, Kono S. Comparison of Japanese, American-Whites and African-Americans--pointers to risk factors to underlying distribution of tumours in the colorectum. *Asian Pac J Cancer Prev*. 2005;6:412-9.