

CURRICULUM VITAE

NAME

Toshiro Ohta, Ph.D.

BIRTH

October, 1965

TITLE

Research Assistant Professor

AFFILIATION

Laboratory of Cell and Molecular Biology of Aging
School of Food and Nutritional Sciences/ Graduate Division of Nutritional and
Environmental Sciences, University of Shizuoka, Japan

EDUCATION

B.S. 1988 University of Tokyo, Faculty of Science
M.S. 1990 University of Tokyo, Graduate School of Science
Ph.D. 1993 University of Tokyo, Graduate School of Science

ACADEMIC CAREER HISTORY

1993-1994 Postdoctoral associate at Japanese Society for Promotion of Science
1994-1996 Postdoctoral associate at University of Minnesota
1996-present Research Assistant Professor at School of Food and Nutritional
Sciences/ Graduate School of Nutritional and Environmental Sciences,
University of Shizuoka

SELECTED PUBLICATIONS

- 1) Kunimasa K, Kobayashi T, Kaji K, Ohta T.
Antiangiogenic effects of indole-3-carbinol and 3,3'-diindolylmethane are associated
with their differential regulation of ERK1/2 and Akt in tube-forming HUVEC.
J Nutr. 2010 Jan;140(1):1-6.
- 2) Ahn M-R, Kunimasa K, Kumazawa S, Nakayama T, Kaji K, Uto Y, Hori H,
Nagasawa H, Ohta T.
Correlation between antiangiogenic activity and antioxidant activity of various
components from propolis.
Mol Nutr Food Res. 2009 May;53(5):643-651.

- 3) Eguchi R, Toné S, Suzuki A, Fujimori Y, Nakano T, Kaji K, Ohta T.
Possible involvement of caspase-6 and -7 but not caspase-3 in the regulation of hypoxia-induced apoptosis in tube-forming endothelial cells.
Exp Cell Res. 2009 Jan 15;315(2):327-335.
- 4) Eguchi R, Naitou H, Kunimasa K, Ayuzawa R, Fujimori Y, Ohashi N, Kaji K, Ohta T.
Proteomic analysis of hypoxia-induced tube breakdown of an in vitro capillary model composed of HUVECs: potential role of p38-regulated reduction of HSP27.
Proteomics. 2008 Jul;8(14):2897-2906.
- 5) Kunimasa K, Kobayashi T, Sugiyama S, Kaji K, Ohta T.
Indole-3-carbinol suppresses tumor-induced angiogenesis by inhibiting tube formation and inducing apoptosis.
Biosci Biotechnol Biochem. 2008 Aug;72(8):2243-2246.
- 6) Ohta T, Kunimasa K, Kobayashi T, Sakamoto M, Kaji K.
Propolis suppresses tumor angiogenesis by inducing apoptosis in tube-forming endothelial cells.
Biosci Biotechnol Biochem. 2008 Sep;72(9):2436-2440.
- 7) Ohta T, Eguchi R, Suzuki A, Miyakaze S, Ayuzawa R, Kaji K.
Hypoxia-induced apoptosis and tube breakdown are regulated by p38 MAPK but not by caspase cascade in an in vitro capillary model composed of human endothelial cells.
J Cell Physiol. 2007 Jun;211(3):673-681.
- 8) Ahn MR, Kunimasa K, Ohta T, Kumazawa S, Kamihira M, Kaji K, Uto Y, Hori H, Nagasawa H, Nakayama T.
Suppression of tumor-induced angiogenesis by Brazilian propolis: Major component artepillin C inhibits in vitro tube formation and endothelial cell proliferation.
Cancer Lett. 2007 Jul 18;252(2):235-243.
- 9) Eguchi R, Suzuki A, Miyakaze S, Kaji K, Ohta T.
Hypoxia induces apoptosis of HUVECs in an in vitro capillary model by activating proapoptotic signal p38 through suppression of ERK1/2.
Cell Signal. 2007 Jun;19(6):1121-1131.
- 10) Ohta T, Essner R, Ryu JH, Palazzo RE, Uetake Y, Kuriyama R.

Characterization of Cep135, a novel coiled-coil centrosomal protein involved in microtubule organization in mammalian cells.
J Cell Biol. 2002 Jan 7;156(1):87-99.

11) Kondo T, Ohta T, Igura K, Hara Y, Kaji K.
Tea catechins inhibit angiogenesis in vitro, measured by human endothelial cell growth, migration and tube formation, through inhibition of VEGF receptor binding.
Cancer Lett. 2002 Jun 28;180(2):139-144.

12) Igura K, Ohta T, Kuroda Y, Kaji K.
Resveratrol and quercetin inhibit angiogenesis in vitro.
Cancer Lett. 2001 Sep 28;171(1):11-16.